arXiv:math/0501159v2 [math.FA] 3 Mar 2005

ON THE CAUCHY–RASSIAS INEQUALITY AND LINEAR n-INNER PRODUCT PRESERVING MAPPINGS

Choonkil Baak[1], Hahng-Yun Chu[2] and Mohammad Sal Moslehian[3]

Abstract. We prove the Cauchy–Rassias stability of linear n-inner product preserving mappings in n-inner product Banach spaces. We apply the Cauchy-Rassias inequality that plays an influencial role in the subject of functional equations. The inequality was introduced for the first time by Th.M.Rassias in his paper entitled: On the stability of the linear mapping in Banach spaces, Proc. Amer. Math.Soc. 72(1978), 297–300.

1. Introduction

Let X and Y be Banach spaces with norms $||\cdot||$ and $\|\cdot\|$, respectively. Consider $f : X \to Y$ to be a mapping such that $f(tx)$ is continuous in $t \in \mathbb{R}$ for each fixed $x \in X$. Th.M. Rassias [23] introduced the following inequality, that we call *Cauchy–Rassias inequality* : Assume that there exist constants $\theta \ge 0$ and $p \in [0, 1)$ such that

$$\|f(x+y) - f(x) - f(y)\| \le \theta(||x||^p + ||y||^p) \tag{*}$$

for all $x, y \in X$. Rassias [23] showed that there exists a unique $\mathbb{R}$-linear mapping $T : X \to Y$ such that

$$\|f(x) - T(x)\| \le \frac{2\theta}{2 - 2^p}||x||^p$$

for all $x \in X$. The inequality $(*)$ has provided a lot of influence in the development of what we now call *Hyers–Ulam–Rassias stability* of functional equations. Beginning around the year 1980 the topic stability of linear functional equations has been studied by a number of mathematicians (see [7], [8] and [11]–[27]).

1991 *Mathematics Subject Classification.* Primary 39B52, 46B04, 46B20, 51Kxx.

Key words and phrases. Cauchy–Rassias stability, linear n-inner product preserving mapping, n-inner product Banach space, Hilbert space.

Supported by Korea Research Foundation Grant KRF-2004-041-C00023.

Typeset by $\mathcal{AMS}$-TEX

Jun and Lee [9] proved the following: Denote by $\varphi : X \setminus \{0\} \times X \setminus \{0\} \to [0, \infty)$ a function such that

$$\widetilde{\varphi}(x, y) = \sum_{j=0}^{\infty} \frac{1}{3^j} \varphi(3^j x, 3^j y) < \infty$$

for all $x, y \in X \setminus \{0\}$. Suppose that $f : X \to Y$ is a mapping satisfying

$$\|2f(\frac{x+y}{2}) - f(x) - f(y)\| \le \varphi(x, y)$$

for all $x, y \in X \setminus \{0\}$. Then there exists a unique additive mapping $T : X \to Y$ such that

$$\|f(x) - f(0) - T(x)\| \le \frac{1}{3}\big(\widetilde{\varphi}(x, -x) + \widetilde{\varphi}(-x, 3x)\big)$$

for all $x \in X \setminus \{0\}$.

Let X and Y be complex Hilbert spaces. A mapping $f : X \to Y$ is called an *inner product preserving mapping* if f satisfies the *orthogonality equation*

$$\langle f(x), f(y) \rangle = \langle x, y \rangle$$

for all $x, y \in X$. The inner product preserving mapping problem has been investigated in several papers (see [2]–[4]).

In this paper, we prove the Cauchy–Rassias stability of inner product preserving mappings in Hilber spaces, introduce the concept of n-inner product Banach space and establish the Cauchy–Rassias stability of linear n-inner product preserving mappings in n-inner product Banach spaces.

2. Cauchy–Rassias stability of inner product preserving mappings in Hilbert spaces

In this section, assume that X is a complex Hilbert space with norm $|| \cdot ||$, and that Y is a complex Hilbert space with norm $\| \cdot \|$.

Theorem 2.1. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exists a function $\varphi : X \times X \to [0, \infty)$ such that*

$$\widetilde{\varphi}(x, y) := \sum_{j=0}^{\infty} \frac{1}{2^j} \varphi(2^j x, 2^j y) < \infty, \tag{2.i}$$

$$\|f(\mu x + \mu y) - \mu f(x) - \mu f(y)\| \le \varphi(x, y), \tag{2.ii}$$

$$| \langle f(x), f(y) \rangle - \langle x, y \rangle | \le \varphi(x, y) \tag{2.iii}$$

for all $\mu \in \mathbb{T}^1 := \{\lambda \in \mathbb{C} \mid |\lambda| = 1\}$ *and all* $x, y \in X$. *Then there exists a unique inner product preserving mapping* $U : X \to Y$ *such that*

$$\|f(x) - U(x)\| \leq \frac{1}{2}\widetilde{\varphi}(x, x) \tag{2.iv}$$

for all $x \in X$.

Proof. By the same reasoning as in the proof of [21, Theorem 2.1], it follows from (2.i) and (2.ii) that there exists a unique $\mathbb{C}$-linear mapping $U : X \to Y$ satisfying (2.iv). The $\mathbb{C}$-linear mapping $U : X \to Y$ is given by

$$U(x) = \lim_{l\to\infty} \frac{1}{2^l} f(2^l x) \tag{2.1}$$

for all $x \in X$.

It follows from (2.iii) that

$$\begin{aligned} \mid \langle \frac{1}{2^l} f(2^l x), \frac{1}{2^l} f(2^l y)\rangle - \langle x, y\rangle \mid &= \frac{1}{2^{2l}} \mid \langle f(2^l x), f(2^l y)\rangle - \langle 2^l x, 2^l y\rangle \mid \\ &\leq \frac{1}{2^{2l}} \varphi(2^l x, 2^l y) \leq \frac{1}{2^l} \varphi(2^l x, 2^l y), \end{aligned}$$

which tends to zero as $l \to \infty$ for all $x, y \in X$ by (2.i). By (2.1),

$$\langle U(x), U(y)\rangle = \lim_{l\to\infty} \langle \frac{1}{2^l} f(2^l x), \frac{1}{2^l} f(2^l y)\rangle = \langle x, y\rangle$$

for all $x, y \in X$, as desired. □

Corollary 2.2. *Let* $f : X \to Y$ *be a mapping with* $f(0) = 0$ *for which there exist constants* $\theta \geq 0$ *and* $p \in [0, 1)$ *such that*

$$\begin{gathered} \|f(\mu x + \mu y) - \mu f(x) - \mu f(y)\| \leq \theta(||x||^p + ||y||^p), \\ \mid \langle f(x), f(y)\rangle - \langle x, y\rangle \mid \leq \theta(||x||^p + ||y||^p) \end{gathered}$$

for all $\mu \in \mathbb{T}^1$ *and all* $x, y \in X$. *Then there exists a unique inner product preserving mapping* $U : X \to Y$ *such that*

$$\|f(x) - U(x)\| \leq \frac{2\theta}{2 - 2^p} ||x||^p$$

for all $x \in X$.

Proof. Define $\varphi(x, y) = \theta(||x||^p + ||y||^p)$ to be Th.M. Rassias upper bound in the Cauchy–Rassias inequality, and apply Theorem 2.1. □

Theorem 2.3. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exists a function $\varphi : X \setminus \{0\} \times X \setminus \{0\} \to [0, \infty)$ such that*

$$\widetilde{\varphi}(x, y) := \sum_{j=0}^{\infty} \frac{1}{3^j} \varphi(3^j x, 3^j y) < \infty, \tag{2.v}$$

$$\|2f(\frac{\mu x + \mu y}{2}) - \mu f(x) - \mu f(y)\| \le \varphi(x, y), \tag{2.vi}$$

$$|\ \langle f(x), f(y)\rangle - \langle x, y\rangle\ | \le \varphi(x, y) \tag{2.vii}$$

for all $\mu \in \mathbb{T}^1$ and all $x, y \in X \setminus \{0\}$. Then there exists a unique inner product preserving mapping $U : X \to Y$ such that

$$\|f(x) - U(x)\| \le \frac{1}{3}\big(\widetilde{\varphi}(x, -x) + \widetilde{\varphi}(-x, 3x)\big) \tag{2.viii}$$

for all $x \in X \setminus \{0\}$.

Proof. By the same reasoning as in the proof of [11, Theorem 2.5], it follows from (2.v) and (2.vi) that there exists a unique $\mathbb{C}$-linear mapping $U : X \to Y$ satisfying (2.viii). The $\mathbb{C}$-linear mapping $U : X \to Y$ is given by

$$U(x) = \lim_{l\to\infty} \frac{1}{3^l} f(3^l x) \tag{2.2}$$

for all $x \in X$.

It follows from (2.vii) that

$$\begin{aligned} |\ \langle \frac{1}{3^l} f(3^l x), \frac{1}{3^l} f(3^l y)\rangle - \langle x, y\rangle\ | &= \frac{1}{3^{2l}} |\ \langle f(3^l x), f(3^l y)\rangle - \langle 3^l x, 3^l y\rangle\ | \\ &\le \frac{1}{3^{2l}} \varphi(3^l x, 3^l y) \le \frac{1}{3^l} \varphi(3^l x, 3^l y), \end{aligned}$$

which tends to zero as $l \to \infty$ for all $x, y \in X$ by (2.v). By (2.2),

$$\langle U(x), U(y)\rangle = \lim_{l\to\infty} \langle \frac{1}{3^l} f(3^l x), \frac{1}{3^l} f(3^l y)\rangle = \langle x, y\rangle$$

for all $x, y \in X$, as desired. □

Corollary 2.4. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exist constants $\theta \ge 0$ and $p \in [0, 1)$ such that*

$$\begin{aligned} \|2f(\frac{\mu x + \mu y}{2}) - \mu f(x) - \mu f(y)\| &\le \theta(||x||^p + ||y||^p), \\ |\ \langle f(x), f(y)\rangle - \langle x, y\rangle\ | &\le \theta(||x||^p + ||y||^p) \end{aligned}$$

for all $\mu \in \mathbb{T}^1$ *and all* $x, y \in X \setminus \{0\}$. *Then there exists a unique inner product preserving mapping* $U : X \to Y$ *such that*

$$\|f(x) - U(x)\| \leq \frac{3+3^p}{3-3^p}\theta||x||^p$$

for all $x \in X \setminus \{0\}$.

Proof. Define $\varphi(x,y) = \theta(||x||^p + ||y||^p)$, and apply Theorem 2.3. □

Theorem 2.5. *Let* $f : X \to Y$ *be a mapping with* $f(0) = 0$ *for which there exists a function* $\varphi : X \setminus \{0\} \times X \setminus \{0\} \to [0, \infty)$ *such that*

$$\sum_{j=0}^{\infty} 3^{2j}\varphi(\frac{x}{3^j}, \frac{y}{3^j}) < \infty, \tag{2.ix}$$

$$\|2f(\frac{\mu x + \mu y}{2}) - \mu f(x) - \mu f(y)\| \leq \varphi(x,y), \tag{2.x}$$

$$| \ \langle f(x), f(y)\rangle - \langle x, y\rangle \ | \leq \varphi(x,y) \tag{2.xi}$$

for all $\mu \in \mathbb{T}^1$ *and all* $x, y \in X \setminus \{0\}$. *Then there exists a unique inner product preserving mapping* $U : X \to Y$ *such that*

$$\|f(x) - U(x)\| \leq \widetilde{\varphi}(\frac{x}{3}, -\frac{x}{3}) + \widetilde{\varphi}(-\frac{x}{3}, x) \tag{2.xii}$$

for all $x \in X \setminus \{0\}$, *where*

$$\widetilde{\varphi}(x,y) := \sum_{j=0}^{\infty} 3^j \varphi(\frac{x}{3^j}, \frac{y}{3^j})$$

for all $x, y \in X \setminus \{0\}$.

Proof. Note that

$$3^j\varphi(\frac{x}{3^j}, \frac{y}{3^j}) \leq 3^{2j}\varphi(\frac{x}{3^j}, \frac{y}{3^j}) \tag{2.3}$$

for all $x, y \in X$ and all positive integers j. By the Jun and Lee's theorem [9, Theorem 6], it follows from (2.ix), (2.3) and (2.x) that there exists a unique additive mapping $U : X \to Y$ satisfying (2.xii). The additive mapping $U : X \to Y$ is given by

$$U(x) = \lim_{l\to\infty} 3^l f(\frac{x}{3^l}) \tag{2.4}$$

for all $x \in X$. By the same method as in the proof of [11, Theorem 2.5], one can show that the additive mapping $U : X \to Y$ is $\mathbb{C}$-linear.

It follows from (2.xi) that

$$| \langle 3^l f(\frac{x}{3^l}), 3^l f(\frac{y}{3^l}) \rangle - \langle x, y \rangle | = 3^{2l} | \langle f(\frac{x}{3^l}), f(\frac{y}{3^l}) \rangle - \langle \frac{x}{3^l}, \frac{y}{3^l} \rangle | \leq 3^{2l} \varphi(\frac{x}{3^l}, \frac{y}{3^l}),$$

which tends to zero as $l \to \infty$ for all $x, y \in X$ by (2.ix). By (2.4),

$$\langle U(x), U(y) \rangle = \lim_{l \to \infty} \langle 3^l f(\frac{x}{3^l}), 3^l f(\frac{y}{3^l}) \rangle = \langle x, y \rangle$$

for all $x, y \in X$, as desired. □

Corollary 2.6. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exist constants $\theta \geq 0$ and $p \in (2, \infty)$ such that*

$$\|2f(\frac{\mu x + \mu y}{2}) - \mu f(x) - \mu f(y)\| \leq \theta(||x||^p + ||y||^p),$$
$$| \langle f(x), f(y) \rangle - \langle x, y \rangle | \leq \theta(||x||^p + ||y||^p)$$

for all $\mu \in \mathbb{T}^1$ and all $x, y \in X \setminus \{0\}$. Then there exists a unique inner product preserving mapping $U : X \to Y$ such that

$$\|f(x) - U(x)\| \leq \frac{3^p + 3}{3^p - 3} \theta ||x||^p$$

for all $x \in X \setminus \{0\}$.

Proof. Define $\varphi(x, y) = \theta(||x||^p + ||y||^p)$, and apply Theorem 2.5. □

3. CAUCHY–RASSIAS STABILITY OF LINEAR n-INNER PRODUCT PRESERVING MAPPINGS IN n-INNER PRODUCT BANACH SPACES

We first recall the notion of n-inner product space.

Definition 3.1 [5]**.** Let X be a complex linear space with $\dim X \geq n \geq 2$ and $\langle \cdot, \cdot \mid \cdot, \cdots, \cdot \rangle : X^{n+1} \to \mathbb{C}$ be a function. Then $(X, \langle \cdot, \cdot \mid \cdot, \cdots, \cdot \rangle)$ is called an *n-inner product space* if

(nI$_1$) $\langle x, x \mid x_2, \cdots, x_n \rangle \geq 0$,

(nI$_2$) $\langle x, x \mid x_2, \cdots, x_n \rangle = 0$if and only if$x, x_2, \cdots, x_n$ are linearly dependent,

(nI$_3$) $\langle x, y \mid x_2, \cdots, x_n \rangle = \overline{\langle y, x \mid x_2, \cdots, x_n \rangle}$,

(nI$_4$) $\langle x, y \mid x_2, \cdots, x_n \rangle = \langle x, y \mid x_{j_2}, \cdots, x_{j_n} \rangle$ for every permutation $(j_2, \cdots j_n)$ of $(2, \cdots, n)$,

(nI$_5$) $\langle x, x \mid x_2, x_3, \cdots, x_n \rangle = \langle x_2, x_2 \mid x, x_3, \cdots, x_n \rangle$,

(nI$_6$) $\langle \alpha x, y \mid x_2, \cdots, x_n \rangle = \alpha \langle x, y \mid x_2, \cdots, x_n \rangle$,

(nI$_7$) $\langle x + y, z \mid x_2, \cdots, x_n \rangle \leq \langle x, z \mid x_2, \cdots, x_n \rangle + \langle y, z \mid x_2, \cdots, x_n \rangle$

for all $\alpha \in \mathbb{C}$ and all $x, y, z, x_1, \cdots, x_n \in X$. The function $\langle \cdot, \cdot \mid \cdot, \cdots, \cdot \rangle$ is called an *n-inner product on X*.

The concept of n-inner product was introduced in [1] for $n = 2$ and in [10] for $n \geq 2$.

For instance, on a given inner product space with inner product $\langle\ ,\ \rangle$, one can put an n-inner product by defining $\langle x, y \mid x_2, \cdots, x_n \rangle$ to be

$$\det \begin{pmatrix} \langle x, y \rangle & \langle x, x_2 \rangle & \cdots & \langle x, x_n \rangle \\ \langle x_2, y \rangle & \langle x_2, x_2 \rangle & \cdots & \langle x_2, x_n \rangle \\ \vdots & \vdots & \cdots & \vdots \\ \langle x_n, y \rangle & \langle x_n, x_2 \rangle & \cdots & \langle x_n, x_n \rangle . \end{pmatrix}$$

Definition 3.2. An n-inner product and normed (respectively, Banach) space X with n-inner product $\langle \cdot, \cdot \mid \cdot, \cdots, \cdot \rangle_X$ and norm $\| \cdot \|$ is called an *n-inner product normed (respectively, Banach) space.*

For example, if $(X, \langle \cdot, \cdot \mid \cdot, \cdots, \cdot \rangle)$ is an n-inner product space, $\{a_1, \cdots, a_n\}$ is a fixed linearly independent set and $k > 0$ then $\langle x, y \rangle := k \sum_{\{i_2, \cdots, i_n\} \subseteq \{1, \cdots, n\}} \langle x, y \mid a_{i_2}, \cdots, a_{i_n} \rangle$ is an inner product on X (see [6]) and so X equipped with the norm induced by this inner product is an n-inner product normed space.

In the rest of this section, assume that X is an n-inner product Banach space with n-inner product $\langle \cdot, \cdot \mid \cdot, \cdots, \cdot \rangle_X$ and norm $\| \cdot \|$, and that Y is an n-inner product Banach space with n-inner product $\langle \cdot, \cdot \mid \cdot, \cdots, \cdot \rangle_Y$ and norm $\| \cdot \|$.

Theorem 3.1. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exists a function $\varphi : X^{n+1} \to [0, \infty)$ such that*

$$\widetilde{\varphi}(x_0, \cdots, x_n) := \sum_{j=0}^{\infty} \frac{1}{2^j} \varphi(2^j x_0, \cdots, 2^j x_n) < \infty, \tag{3.i}$$

$$\| f(\mu x_0 + \mu x_1) - \mu f(x_0) - \mu f(x_1) \| \leq \varphi(x_0, x_1, \underbrace{0, \cdots, 0}_{n-1 \text{ times}}), \tag{3.ii}$$

$$\begin{aligned} &| \langle f(x_0), f(x_1) \mid f(x_2), \cdots, f(x_n) \rangle_Y - \langle x_0, x_1 \mid x_2, \cdots, x_n \rangle_X | \\ &\qquad \leq \varphi(x_0, \cdots, x_n) \end{aligned} \tag{3.iii}$$

for all $\mu \in \mathbb{T}^1$ and all $x_0, \cdots, x_n \in X$. Then there exists a unique $\mathbb{C}$-linear n-inner product preserving mapping $U : X \to Y$ such that

$$\| f(x) - U(x) \| \leq \frac{1}{2} \widetilde{\varphi}(x, x, \underbrace{0, \cdots, 0}_{n-1 \text{ times}}) \tag{3.iv}$$

for all $x \in X$.

Proof. By the same reasoning as in the proof of [20, Theorem 2.1], it follows from (3.i) and (3.ii) that there exists a unique $\mathbb{C}$-linear mapping $U : X \to Y$ satisfying (3.iv). The $\mathbb{C}$-linear mapping $U : X \to Y$ is given by

$$U(x) = \lim_{l\to\infty} \frac{1}{2^l} f(2^l x) \tag{3.1}$$

for all $x \in X$.

It follows from (3.iii) that

$$\begin{aligned} &| \langle \frac{1}{2^l} f(2^l x_0), \frac{1}{2^l} f(2^l x_1) \, | \frac{1}{2^l} f(2^l x_2), \cdots, \frac{1}{2^l} f(2^l x_n)\rangle_Y \\ &\qquad - \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \mid \\ &\qquad = \frac{1}{2^{2nl}} | \, \langle f(2^l x_0), f(2^l x_1) \mid f(2^l x_2), \cdots, f(2^l x_n)\rangle_Y \\ &\qquad\qquad - \langle 2^l x_0, 2^l x_1 \mid 2^l x_2, \cdots, 2^l x_n\rangle_X \mid \\ &\qquad \le \frac{1}{2^{2nl}} \varphi(2^l x_0, \cdots, 2^l x_n) \le \frac{1}{2^l} \varphi(2^l x_0, \cdots, 2^l x_n), \end{aligned}$$

which tends to zero as $l \to \infty$ for all $x_0, \cdots, x_n \in X$ by (3.i). By (3.1),

$$\begin{aligned} &\langle U(x_0), U(x_1) \mid U(x_2), \cdots, U(x_n)\rangle_Y \\ &\qquad = \lim_{l\to\infty} \langle \frac{1}{2^l} f(2^l x_0), \frac{1}{2^l} f(2^l x_1) \mid \frac{1}{2^l} f(2^l x_2), \cdots, \frac{1}{2^l} f(2^l x_n)\rangle_Y \\ &\qquad = \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \end{aligned}$$

for all $x_0, \cdots, x_n \in X$, as desired. $\square$

Corollary 3.2. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exist constants $\theta \ge 0$ and $p \in [0, 1)$ such that*

$$\|f(\mu x_0 + \mu x_1) - \mu f(x_0) - \mu f(x_1)\| \le \theta(||x_0||^p + ||x_1||^p),$$

$$| \, \langle f(x_0), f(x_1) \mid f(x_2), \cdots, f(x_n)\rangle_Y - \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \mid \le \theta \sum_{j=0}^{n} ||x_j||^p$$

for all $\mu \in \mathbb{T}^1$ and all $x_0, \cdots, x_n \in X$. Then there exists a unique $\mathbb{C}$-linear n-inner product preserving mapping $U : X \to Y$ such that

$$\|f(x) - U(x)\| \le \frac{2\theta}{2 - 2^p} ||x||^p$$

for all $x \in X$.

Proof. Define $\varphi(x_0, \cdots, x_n) = \theta \sum_{j=0}^{n} ||x_j||^p$ to be Th.M. Rassias upper bound in the Cauchy–Rassias inequality, and apply Theorem 3.1. $\square$

Theorem 3.3. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exists a function $\varphi : (X \setminus \{0\})^{n+1} \to [0, \infty)$ such that*

$$\widetilde{\varphi}(x_0, \cdots, x_n) := \sum_{j=0}^{\infty} \frac{1}{3^j} \varphi(3^j x_0, \cdots, 3^j x_n) < \infty, \tag{3.v}$$

$$\|2f(\frac{\mu x_0 + \mu x_1}{2}) - \mu f(x_0) - \mu f(x_1)\| \le \varphi(x_0, x_1, \underbrace{0, \cdots, 0}_{n-1 \text{ times}}), \tag{3.vi}$$

$$\begin{aligned} &| \langle f(x_0), f(x_1) \mid f(x_2), \cdots, f(x_n)\rangle_Y - \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \mid \\ &\qquad \le \varphi(x_0, \cdots, x_n) \end{aligned} \tag{3.vii}$$

for all $\mu \in \mathbb{T}^1$ and all $x_0, \cdots, x_n \in X \setminus \{0\}$. Then there exists a unique $\mathbb{C}$-linear n-inner product preserving mapping $U : X \to Y$ such that

$$\|f(x) - U(x)\| \le \frac{1}{3}\big(\widetilde{\varphi}(x, -x, \underbrace{0, \cdots, 0}_{n-1 \text{ times}}) + \widetilde{\varphi}(-x, 3x, \underbrace{0, \cdots, 0}_{n-1 \text{ times}})\big) \tag{3.viii}$$

for all $x \in X \setminus \{0\}$.

Proof. By the same reasoning as in the proof of [20, Theorem 2.5], it follows from (3.v) and (3.vi) that there exists a unique $\mathbb{C}$-linear mapping $U : X \to Y$ satisfying (3.viii). The $\mathbb{C}$-linear mapping $U : X \to Y$ is given by

$$U(x) = \lim_{l \to \infty} \frac{1}{3^l} f(3^l x) \tag{3.2}$$

for all $x \in X$.

It follows from (3.vii) that

$$\begin{aligned} | \langle \frac{1}{3^l} f(3^l x_0), \frac{1}{3^l} f(3^l x_1) \mid & \frac{1}{3^l} f(3^l x_2), \cdots, \frac{1}{3^l} f(3^l x_n)\rangle_Y \\ & - \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \mid \\ & = \frac{1}{3^{2nl}} | \langle f(3^l x_0), f(3^l x_1) \mid f(3^l x_2), \cdots, f(3^l x_n)\rangle_Y \\ & \qquad - \langle 3^l x_0, 3^l x_1 \mid 3^l x_2, \cdots, 3^l x_n\rangle_X \mid \\ & \le \frac{1}{3^{2nl}} \varphi(3^l x_0, \cdots, 3^l x_n) \le \frac{1}{3^l} \varphi(3^l x_0, \cdots, 3^l x_n), \end{aligned}$$

which tends to zero as $l \to \infty$ for all $x_0, \cdots, x_n \in X$ by (3.v). By (3.2),

$$\begin{aligned} &\langle U(x_0), U(x_1) \mid U(x_2), \cdots, U(x_n)\rangle_Y \\ &\qquad = \lim_{l \to \infty} \langle \frac{1}{3^l} f(3^l x_0), \frac{1}{3^l} f(3^l x_1) \mid \frac{1}{3^l} f(3^l x_2), \cdots, \frac{1}{3^l} f(3^l x_n)\rangle_Y \\ &\qquad = \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \end{aligned}$$

for all $x_0, \cdots, x_n \in X$, as desired. □

Corollary 3.4. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exist constants $\theta \geq 0$ and $p \in [0, 1)$ such that*

$$\|2f(\frac{\mu x_0 + \mu x_1}{2}) - \mu f(x_0) - \mu f(x_1)\| \leq \theta(||x_0||^p + ||x_1||^p),$$

$$| \langle f(x_0), f(x_1) \mid f(x_2), \cdots, f(x_n)\rangle_Y - \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \mid \leq \theta \sum_{j=0}^{n} ||x_j||^p$$

for all $\mu \in \mathbb{T}^1$ and all $x_0, \cdots, x_n \in X \setminus \{0\}$. Then there exists a unique $\mathbb{C}$-linear n-inner product preserving mapping $U : X \to Y$ such that

$$\|f(x) - U(x)\| \leq \frac{3 + 3^p}{3 - 3^p}\theta||x||^p$$

for all $x \in X$.

Proof. Define $\varphi(x_0, \cdots, x_n) = \theta \sum_{j=0}^{n} ||x_j||^p$, and apply Theorem 3.3. □

Theorem 3.5. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exists a function $\varphi : (X \setminus \{0\})^{n+1} \to [0, \infty)$ such that*

$$\sum_{j=0}^{\infty} 3^{2nj}\varphi(\frac{x_0}{3^j}, \cdots, \frac{x_n}{3^j}) < \infty, \tag{3.ix}$$

$$\|2f(\frac{\mu x_0 + \mu x_1}{2}) - \mu f(x_0) - \mu f(x_1)\| \leq \varphi(x_0, x_1, \underbrace{0, \cdots, 0}_{n-1 \text{ times}}), \tag{3.x}$$

$$\begin{aligned} | \langle f(x_0), f(x_1) \mid f(x_2), \cdots, f(x_n)\rangle_Y - \langle x_0, x_1 \mid x_2, \cdots, x_n\rangle_X \mid \\ \leq \varphi(x_0, \cdots, x_n) \end{aligned} \tag{3.xi}$$

for all $\mu \in \mathbb{T}^1$ and all $x_0, \cdots, x_n \in X \setminus \{0\}$. Then there exists a unique $\mathbb{C}$-linear n-inner product preserving mapping $U : X \to Y$ such that

$$\|f(x) - U(x)\| \leq \widetilde{\varphi}(\frac{x}{3}, -\frac{x}{3}, \underbrace{0, \cdots, 0}_{n-1 \text{ times}}) + \widetilde{\varphi}(-\frac{x}{3}, x, \underbrace{0, \cdots, 0}_{n-1 \text{ times}}) \tag{3.xii}$$

for all $x \in X \setminus \{0\}$, where

$$\widetilde{\varphi}(x_0, \cdots, x_n) := \sum_{j=0}^{\infty} 3^j \varphi(\frac{x_0}{3^j}, \cdots, \frac{x_n}{3^j})$$

for all $x_0, \cdots, x_n \in X$.

Proof. Note that

$$3^j\varphi(\frac{x_0}{3^j},\cdots,\frac{x_n}{3^j}) \le 3^{2nj}\varphi(\frac{x_0}{3^j},\cdots,\frac{x_n}{3^j}) \tag{3.3}$$

for all $x_0,\cdots,x_n \in X$ and all positive integers j. By the same reasoning as in the proof of Theorem 2.5, it follows from (3.ix), (3.3) and (3.x) that there exists a unique $\mathbb{C}$-linear mapping $U: X \to Y$ satisfying (3.xii). The $\mathbb{C}$-linear mapping $U: X \to Y$ is given by

$$U(x) = \lim_{l\to\infty} 3^l f(\frac{x}{3^l}) \tag{3.4}$$

for all $x \in X$.

It follows from (3.xi) that

$$\begin{aligned}
\mid \langle 3^l f(\frac{x_0}{3^l}), 3^l f(\frac{x_1}{3^l}) \mid 3^l f(\frac{x_2}{3^l}),\cdots, 3^l f(\frac{x_n}{3^l})\rangle_Y & \\
- \langle x_0, x_1 \mid x_2,\cdots,x_n\rangle_X \mid & \\
= 3^{2nl}\mid \langle f(\frac{x_0}{3^l}), f(\frac{x_1}{3^l}) \mid f(\frac{x_2}{3^l}),\cdots, f(\frac{x_n}{3^l})\rangle_Y & \\
- \langle \frac{x_0}{3^l}, \frac{x_1}{3^l} \mid \frac{x_2}{3^l},\cdots,\frac{x_n}{3^l}\rangle_X \mid & \\
\le 3^{2nl}\varphi(\frac{x_0}{3^l},\cdots,\frac{x_n}{3^l}), &
\end{aligned}$$

which tends to zero as $l \to \infty$ for all $x_0,\cdots,x_n \in X$ by (3.ix). By (3.4),

$$\begin{aligned}
&\langle U(x_0), U(x_1) \mid U(x_2),\cdots,U(x_n)\rangle_Y \\
&\quad = \lim_{l\to\infty}\langle 3^l f(\frac{x_0}{3^l}), 3^l f(\frac{x_1}{3^l}) \mid 3^l f(\frac{x_2}{3^l}),\cdots, 3^l f(\frac{x_n}{3^l})\rangle_Y \\
&\quad = \langle x_0, x_1 \mid x_2,\cdots,x_n\rangle_X
\end{aligned}$$

for all $x_0,\cdots,x_n \in X$, as desired. □

Corollary 3.6. *Let $f : X \to Y$ be a mapping with $f(0) = 0$ for which there exist constants $\theta \ge 0$ and $p \in (2n,\infty)$ such that*

$$\|2f(\frac{\mu x_0 + \mu x_1}{2}) - \mu f(x_0) - \mu f(x_1)\| \le \theta(||x_0||^p + ||x_1||^p),$$

$$\mid \langle f(x_0), f(x_1) \mid f(x_2),\cdots,f(x_n)\rangle_Y - \langle x_0, x_1 \mid x_2,\cdots,x_n\rangle_X \mid \le \theta\sum_{j=0}^{n}||x_j||^p$$

for all $\mu \in \mathbb{T}^1$ and all $x_0,\cdots,x_n \in X\setminus\{0\}$. Then there exists a unique $\mathbb{C}$-linear n-inner product preserving mapping $U : X \to Y$ such that

$$\|f(x) - U(x)\| \le \frac{3^p+3}{3^p-3}\theta||x||^p$$

for all $x \in X \setminus \{0\}$.

Proof. Define $\varphi(x_0, \cdots, x_n) = \theta \sum_{j=0}^{n} ||x_j||^p$, and apply Theorem 3.5. $\square$

Acknowledgment. We sincerely thank Professor Themistocles M. Rassias for his valuable suggestions.

References

1. C. Diminnie, S. Gähler and A. White, 2-*inner product spaces*, Demonstratio Math. **6** (1973), 525–536.
2. J. Chmieliński, *On a singular case in the Hyers–Ulam–Rassias stability of the Wigner equation*, J. Math. Anal. Appl. **289** (2004), 571–583.
3. ______, *Linear mappings approximately preserving orthogonality*, J. Math. Anal. Appl. (to appear).
4. J. Chmieliński and S. Jung, *The stability of the Wigner equation on a restricted domain*, J. Math. Anal. Appl. **254** (2001), 309–320.
5. Y.J. Cho, P.C.S. Lin, S.S. Kim and A. Misiak, *Theory of* 2-*Inner Product Spaces*, Nova Science Publ., New York, 2001.
6. H. Gunawan, *An inner product space that makes a set of vectors orthonormal*, J. Austral. Math. Soc. Gaz. **28** (2001), 194–197.
7. D.H. Hyers, G. Isac and Th.M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhauser, Boston, Basel, Berlin, 1998.
8. D.H. Hyers and Th.M. Rassias, *Approximate homomorphisms*, Aequationes Mathematicae **44** (1992), 125–153.
9. K. Jun and Y. Lee, *A generalization of the Hyers–Ulam–Rassias stability of Jensen's equation*, J. Math. Anal. Appl. **238** (1999), 305–315.
10. A. Misiak, *n-inner product spaces*, Math. Nachr. **140** (1989), 299–319.
11. C. Park, *On the stability of the linear mapping in Banach modules*, J. Math. Anal. Appl. **275** (2002), 711–720.
12. ______, *Linear functional equations in Banach modules over a* C^**-algebra*, Acta Appl. Math. **77** (2003), 125–161.
13. ______, *Modified Trif's functional equations in Banach modules over a* C^**-algebra and approximate algebra homomorphisms*, J. Math. Anal. Appl. **278** (2003), 93–108.
14. ______, *Lie* $*$*-homomorphisms between Lie* C^**-algebras and Lie* $*$*-derivations on Lie* C^**-algebras*, J. Math. Anal. Appl. **293** (2004), 419–434.
15. ______, *Universal Jensen's equations in Banach modules over a* C^**-algebra and its unitary group*, Acta Math. Sinica **20** (2004), 1047–1056.
16. ______, *Homomorphisms between Lie* JC^**-algebras and Cauchy–Rassias stability of Lie* JC^**-algebra derivations*, J. Lie Theory **15** (2005), 393–414.
17. ______, *Isomorphisms between unital* C^**-algebras*, Math. Ineq. Appl. (to appear).
18. ______, *Positive linear operators in* C^**-algebras*, Math. Ineq. Appl. (to appear).
19. ______, *Homomorphisms between* C^**-algebras and linear derivations on* C^**-algebras*, Math. Ineq. Appl. (to appear).
20. ______, *Homomorphisms between Poisson* JC^**-algebras*, Bull. Braz. Math. Soc. (to appear).
21. C. Park and J. Hou, *Homomorphisms between* C^**-algebras associated with the Trif functional equation and linear derivations on* C^**-algebras*, J. Korean Math. Soc. **41** (2004), 461–477.
22. A. Prastaro and Th.M. Rassias, *Ulam stability in geometry of PDE's*, Nonlinear Functional Analysis and Applications **8** (2003), 259–278.
23. Th.M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.

24. ______, *The problem of S.M. Ulam for approximately multiplicative mappings*, J. Math. Anal. Appl. **246** (2000), 352–378.
25. ______, *On the stability of functional equations in Banach spaces*, J. Math. Anal. Appl. **251** (2000), 264–284.
26. ______, *On the stability of functional equations and a problem of Ulam*, Acta Applicandae Mathematicae **62** (2000), 23–130.
27. Th.M. Rassias and P. Šemrl, *On the Hyers–Ulam stability of linear mappings*, J. Math. Anal. Appl. **173** (1993), 325–338.

[1,2]Department of Mathematics, Chungnam National University, Daejeon 305–764, South Korea; [3]Department of Mathematics, Ferdowsi University, P.O. Box 1159, Mashhad 91775, Iran

E-mail address: [1]`cgpark@cnu.ac.kr`; [2]`hychu@math.cnu.ac.kr`; [3]`msalm@math.um.ac.ir`